

PREMISE TREATMENT



Fly and Roach Control

- ✓ Kills Pit & Filth Flies, Mosquitoes, and Cockroaches
- ✓ Ready to use
- ✓ Contact insecticide
- ✓ Safe for use around people and animals when used as directed

- ✓ For use on livestock premises, including: swine quarters, pens and shelters



NET CONTENTS: 2.5 gallons (9.46 L)

Active Ingredients:

Geraniol.....	1.36%
Cinnamon Oil.....	0.20%
Inert ingredients*	98.44%
TOTAL.....	100.0%

*White mineral oil, isopropyl myristate, butyl lactate, triethyl citrate, vanillin
Contains petroleum distillates

Keep Out of Reach of Children

CAUTION: See Precautionary Statements



Manufactured by:



5151 McCrimmon Parkway, Suite 275
Morrisville, NC 27560

PureScience and TyraTech logo are trademarks of TyraTech, Inc.

Patents: tyratech.com/legal

This product has not been registered by the United States Environmental Protection Agency. TyraTech, Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act.

40350FRT2018200

First Aid

- **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- **IF ON SKIN:** Wash thoroughly with soap and water. If irritation develops, contact a doctor or poison control center for treatment advice.
- **IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
- **IF INHALED:** Move affected person(s) to fresh air. Call a doctor for further advice if there is difficulty breathing.

Have the product label or container with you when seeking treatment or medical advice.

Note to Physicians

Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

Precautionary Statements

Hazards to Humans

CAUTION: Causes serious eye irritation. Prolonged or frequent, repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE): Appropriate personal protection equipment should be used where and when applicable. Applicators should wear protective gloves, clothing, eye protection, face protection.

User Safety Recommendations: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Avoid contact with eyes, skin, clothing, or plants. Avoid breathing spray. Wash contaminated clothing separately.

Physical & Chemical Hazards

Do not apply on or around electrical wiring. Do not apply to equipment when hot. Treated surfaces may be slippery when wet. Excessive residue may be wiped up with a clean cloth.

Use Restrictions

- Apply this product only as specified on this label.
- Only protected applicators may be in treatment area during application.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift.
- Product was designed for premise and spatial application around and over animals where animals may be contacted through drift. Product was not designed to treat ectoparasites on animals.
- **DO NOT** remain in treated area. Wait until product has settled to return to the area. It is not necessary to remove animals from the area being treated.

Directions for Use

Product is ready to use – **DO NOT DILUTE.**

For use wherever insect control is needed, including swine quarters, pens and shelters, and other associated areas. Product use permitted in other areas or facilities when product is used according to label directions.

Application for Adult Flying Insects

METHOD OF APPLICATION: Aerial mist powered by air compressor (e.g. Stihl SR450)

RATE:

- Apply 2.25 – 4.5 ft oz / 1,000 cubic feet of space to be treated
- Apply full strength to inside and outside areas of production facilities to knock down adult populations
- For inside use make every attempt to fill the room being treated with a fine mist or fog to insure good control

TIMING OF APPLICATION: Apply as needed to hold populations at low levels. For example, apply 3 applications at 3-5 day intervals, in order to break the breeding cycle.

Application Procedures for Immature Flying Insects (Farrowing, Gestation, and Finishing):

For use in premises that are not regularly flushed.

METHOD OF APPLICATION: Pressurized backpack or handheld pump sprayer (e.g. Hudson SP2 Backpack sprayer)

RATE:

- Apply 2.8 – 3.5 gallon / 1,000 sq ft of space to be treated
- Apply full strength to any area where standing water occurs, example pits under pigs
- For pits pour 0.5 gallon into the pit every 8 to 10 feet
- Apply to edges of pits and areas with high organic slurry using a pressurized pump sprayer

TIMING OF APPLICATION: Apply as needed to hold populations at low levels. For example, apply 3 applications at 3-5 day intervals, in order to break the breeding cycle.

Types of Flies Controlled

PIT FLIES Phorid and psychodidae

FILTH FLIES House flies

For Cockroaches

Apply product using a pneumatic-pump sprayer. Treat areas inside and outside of production facilities, where cockroaches congregate, targeting harborage and spaces where insects may be present such as ceiling joists, walls, floors, and around pipes or other structural features. Spray to the point of saturation, contacting as many insects as possible.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store at room temperature in a dry area away from heat or open flame. **CONTAINER HANDLING:** Do not reuse this container. **IF EMPTY:** When container is empty, rinse and recycle if available or dispose of in sanitary landfill. **IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

